

University of Groningen

Microspectrofluorometry on Fly Photoreceptors In Vivo. Spectral Properties of Visual Pigment Fluorescence

Kruizinga, B.; Stavenga, D.G.

Published in:
Vision Research

DOI:
[10.1016/0042-6989\(84\)90352-3](https://doi.org/10.1016/0042-6989(84)90352-3)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1984

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kruizinga, B., & Stavenga, D. G. (1984). Microspectrofluorometry on Fly Photoreceptors In Vivo. Spectral Properties of Visual Pigment Fluorescence. *Vision Research*, 24(11), 1699-1699.
[https://doi.org/10.1016/0042-6989\(84\)90352-3](https://doi.org/10.1016/0042-6989(84)90352-3)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

RHODOPSIN-PORPHYROPSIN SYSTEM IN CRAYFISH

TATSUO SUZUKI and MOMOYO MAKINO-TASAKA

Department of Pharmacology, Hyogo College of Medicine, Nishinomiya, Hyogo 663, Japan

A sensitive analytical method was developed to study the rhodopsin-porphyrpsin system in the eye. Oximes of 11-*cis*-retinal, 11-*cis*-3-dehydroretinal, all-*trans*-retinal, and all-*trans*-3-dehydroretinal were quantitatively determined by high-pressure liquid chromatography. The rhodopsin and porphyropsin of bullfrog retina were analyzed with this method. The results agreed with those obtained from the bleaching kinetics of visual pigments extracted with detergent. The analytical method was so sensitive that a 1/500 part of a bullfrog retina was sufficient for analysis. We studied the chromophore of the visual pigment of freshwater crustacean with this analytical method. The dark-adapted eye of crayfish (*Procambarus clarkii*) contained 11-*cis*-3-dehydroretinal which was isomerized to all-*trans* form by light-adaptation. All-*trans*-isomers produced by light were slowly reconverted to 11-*cis*-isomers in the dark with a half-time of 20 h (20°C). Seasonal variation was observed in the 3-dehydroretinal content. The value was below 5% in summer and about 40% in winter. The results suggest that the crayfish has a rhodopsin-porphyropsin system similar to that in freshwater fish.

LOCALIZATION OF DIFFERENT VISUAL PIGMENTS WITHIN THE RHABDOMS OF INSECT COMPOUND EYES

HELMUT LANGER

Institute for Animal Physiology, Ruhr-University, P.O. Box 10 21 48, D-4630 Bochum 1, F.R.G.

After identification of visual pigments in extracts from insect eyes, there remains the problem of their distribution between the different types of visual cells within the ommatidium. Compound eyes after illumination with monochromatic light were investigated by microspectrophotometry and electron microscopy. Photometrically, difference spectra could be obtained from single rhabdomeres of "open" rhabdoms, indicating only one receptor pigment within a rhabdomere. In fused rhabdoms, distinct changes of the microvillar structure were found in the rhabdomeres of different receptor cells after different monochromatic illuminations. The changes were proved to occur *in vivo* only temporarily. Examples presented are the eyes of hemiptera, especially of the pond skater, *Gerris lacustris*, with three pigments, and of nocturnal lepidoptera, especially of the armyworm moth, *Spodoptera exempta*, with four pigments.

This work was done in collaboration with Drs B. Hamann, C. C. Meinecke, P. Schlecht, B. Welsch (Bochum), and Dr R. Schwind (University of Regensburg). It was financially supported by Deutsche Forschungsgemeinschaft.

MICROSPECTROFLUOROMETRY ON FLY PHOTORECEPTORS *IN VIVO*. SPECTRAL PROPERTIES OF VISUAL PIGMENT FLUORESCENCE

B. KRUIZINGA and D. G. STAVENGA

Biophysics Department, Laboratorium voor Algemene Natuurkunde, Rijksuniversiteit Groningen,
The Netherlands

Visual pigment fluorescence has been studied in the compound eyes of houseflies *Musca* and blowflies *Calliphora*, white eyed mutants, *in vivo*. Rhodopsin R fluorescence is negligible, but the metarhodopsin state M fluoresces distinctly in the red. At extreme illumination intensities a 3–5 times stronger fluorescing M' state is populated. Both M and M' have emission spectra, peaking at about 660 nm, which do not noticeably change in the physiological temperature range, so, indicating the rigidity of the visual molecule in the membrane. The fluorescence life times of M and M', measured by applying a nitrogen pulse laser, are shorter than 40 nsec. Metarhodopsin M fluorescence has been used in a study of the photochemical cycle. The results were in good agreement with those obtained by transmission microspectrophotometry [Kruijinga *et al.* (1983), *Biophys. Struct. Mech.* **9**, 299]. Microspectrofluorometry hence can be utilised for analysing photopigment processes in completely intact, living animals.